

AD-A099 821

CORNELL UNIV ITHACA N Y SCHOOL OF OPERATIONS RESEARC--ETC F/G 12/1
CONDITIONS UNDER WHICH E(N1) = INFINITY FOR TONG'S ADAPTIVE SOL--ETC(U)
SEP 80 C E MCCULLOCH DAA629-80-C-0036

UNCLASSIFIED


TR-480

NL

1 of 1
AL A
12/1/80



END
DATE
FILMED
6-81
DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A099821

II

SCHOOL
OF
OPERATIONS RESEARCH
AND
INDUSTRIAL ENGINEERING

See 473



COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK 1485

DTIC
ELECTE
AND NO
KA

SCHOOL OF OPERATIONS RESEARCH
AND INDUSTRIAL ENGINEERING
COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK

TECHNICAL REPORT NO. 480

September 1980

CONDITIONS UNDER WHICH $E\{N_1\} = \infty$

FOR TONG'S ADAPTIVE SOLUTION
TO RANKING AND SELECTION PROBLEMS

by

Charles E. McCulloch
Cornell University

Accession For	
NTIS GRA&I	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special

Research supported by U.S. Army Research Office-Durham Contract DAAG-29-80-C-0036,
Office of Naval Research Contract N00014-75-C-0586 at Cornell University.

Approved for Public Release; Distribution Unlimited.

THE FINDINGS IN THIS REPORT ARE NOT TO BE
CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE
ARMY POSITION, UNLESS SO DESIGNATED BY OTHER
AUTHORIZED DOCUMENTS.

Abstract

Tong [9] proposed an adaptive approach as an alternative to the classical indifference-zone formulation of the problems of ranking and selection. With a fixed pre-selected γ^* ($1/k < \gamma^* < 1$) his procedure calls for the termination of vector-at-a-time sampling when the estimated probability of a correct selection exceeds γ^* for the first time. The purpose of this note is to show that for the case of two normal populations with common known variance, the expected number of vector-observations required by Tong's procedure to terminate sampling approaches infinity as the two population means approach equality for $\gamma^* \geq \sim 0.8413$. This phenomenon presumably also persists if the two largest of $k \geq 3$ population means approach equality. Since in the typical ranking and selection setting it usually is assumed that the experimenter has no knowledge concerning the differences between the population means, the experimenter who uses Tong's procedure clearly does so at his own risk.

1. Introduction. We retain the notation of Tong [9] and consider the important case of Section 5 of his paper, namely that of the normal family. Here

$\{X_{ij}\}_{j=1}^{\infty}$ are i.i.d. normal r.v.'s with means θ_i ($1 \leq i \leq k$) and common known variance σ^2 . For every n ($n = 1, 2, \dots$) let $\bar{X}_i^{(n)} = \frac{1}{n} \sum_{j=1}^n X_{ij}$ ($1 \leq i \leq k$), and denote the ordered sample means by $\bar{X}_{[1]}^{(n)} < \dots < \bar{X}_{[k]}^{(n)}$. Define $\hat{\delta}_i = \hat{\delta}_i^{(n)} = \bar{X}_{[k]}^{(n)} - \bar{X}_{[i]}^{(n)}$ ($1 \leq i \leq k-1$), and

$$\hat{\gamma}_1^{(n)} = \int_{-\infty}^{\infty} \prod_{i=1}^{k-1} \Phi(y + n^{1/2} \hat{\delta}_i / \sigma) d\Phi(y), \quad (1)$$

the latter being an estimate of the probability of a correct selection (PCS) at stage n . (See Gibbons, Olkin, and Sobel [5], Section 2.3.4.) Here $\Phi(\cdot)$ is the standard normal cdf. Tong's primary procedure, R_1 , stops at the first stage that $\hat{\gamma}_1^{(n)} \geq \gamma^*$ where γ^* ($1/k < \gamma^* < 1$) is a pre-selected constant, analogous to P^* in the classical indifference-zone formulation of Bechhofer [1]. Thus the stopping stage for R_1 is $N_1 = \inf\{n: \hat{\gamma}_1^{(n)} \geq \gamma^*\}$ where the infimum of the empty set is defined to be $+\infty$.

Denoting the ordered values of the θ_i by $\theta_{[1]} \leq \dots \leq \theta_{[k]}$, Tong [9] proved for arbitrary $\theta = (\theta_1, \dots, \theta_k)$ with $\theta_{[k-1]} < \theta_{[k]}$ that asymptotically ($\gamma^* \rightarrow 1$) we have $E_{\theta}\{N_1\} \sim n_1^*$ where $n_1^* = n_1^*(\theta)$ is the smallest single-stage sample size that achieves a PCS of γ^* for that θ . If θ' is more (less) favorable to the experimenter than θ , and n_1^* was chosen under the assumption that θ is the true state of nature, then one would anticipate that $E_{\theta'}\{N_1\} < n_1^* (> n_1^*)$. In particular, when $\theta'_{[k-1]} = \theta'_{[k]}$ one might anticipate that $E_{\theta'}\{N_1\} \gg n_1^*$. In Section 2 we prove for $k = 2$ that $E_{\theta}\{N_1\}$ is actually infinite when $\theta_{[1]} = \theta_{[2]}$ and γ^* is moderately large.

2. The case of $k = 2$ normal populations

Our result is summarized in the following theorem:

Theorem: For $k = 2$ and $\theta_{[1]} = \theta_{[2]}$, the r.v. $N_1 = \inf\{n: \hat{\gamma}_1^{(n)} \geq \gamma^*\}$ has $E_{\theta} \{N_1\} = \infty$ when $\phi^{-1}(\gamma^*) \geq 1$.

Remark 2.1: $\phi^{-1}(\gamma^*) \geq 1$ for $\gamma^* \geq \phi(1) \sim 0.8413$. Hence, for $\gamma^* \geq \phi(1)$ we have $E_{\theta} \{N_1\} \rightarrow \infty$ as $\theta_{[2]} - \theta_{[1]} \rightarrow 0$.

Proof: For $k = 2$ normal populations we can write (1) as

$$\begin{aligned} \hat{\gamma}_1^{(n)} &= \int_{-\infty}^{\infty} \phi(y + n^{1/2} \hat{\delta}_1 / \sigma) d\phi(y) \\ &= \phi\left(\sqrt{\frac{n}{2}} \frac{\hat{\delta}_1}{\sigma}\right). \end{aligned} \quad (2)$$

Then $N_1 = \inf\{n: \hat{\delta}_1 \geq \sqrt{\frac{2\sigma^2}{n}} \phi^{-1}(\gamma^*)\}$. Since $\hat{\delta}_1 = |\bar{X}_1^{(n)} - \bar{X}_2^{(n)}|$ we see that

$$\begin{aligned} N_1 &= \inf\{n: \left| \sum_{j=1}^n (X_{1j} - X_{2j}) / \sqrt{2}\sigma \right| \geq n^{1/2} \phi^{-1}(\gamma^*)\} \\ &= \inf\{n: |S_n| \geq n^{1/2} \phi^{-1}(\gamma^*)\} \end{aligned}$$

where for $\theta_1 = \theta_2$ we have that S_n is the sum of n i.i.d. $N(0,1)$ r.v.'s.

By Corollary 1 of Chow, Robbins and Teicher [3], p. 792, we see that $E\{N_1\}$ is infinite for $\phi^{-1}(\gamma^*) \geq 1$, i.e., for $\gamma^* \geq \phi(1) \sim 0.8413$, and thus the theorem is proved.

Remark 2.2: The same result holds for Tong's procedure R_2 since R_2 stops after R_1 .

3. Comparisons of $E\{N_1\}$ with the corresponding quantity for other competing procedures

In the setting of Section 1 with $k \geq 2$, the expected number of stages required to terminate experimentation for three other sequential procedures are compared with the corresponding quantity for Tong's R_1 . The procedures are the closed sequential procedure (with elimination) of Paulson [6] (with the improvement of Fabian [4]), the open sequential procedure (without elimination) of Bechhofer, Kiefer, and Sobel [2], p. 264, and the two-stage procedure (with elimination) of Tamhane and Bechhofer [7], [8]; all of these procedures are known to guarantee the indifference-zone probability requirement for all $k \geq 2$ and specified $\{\delta^*, P^*\}$ ($0 < \delta^* < \infty$, $1/k < P^* < 1$). For Paulson's procedure the number of stages is bounded. For the Bechhofer-Kiefer-Sobel procedure the expected number of stages for $k = 2$ has been proven to be bounded ([2], p. 224); for this procedure with $k \geq 2$ and $\theta_{[1]} = \theta_{[k]}$, an excellent approximation ($P^* \rightarrow 1$) supported by Monte Carlo sampling results is available for the expected number of stages ([2], pp. 296-7). For the Tamhane-Bechhofer procedure the number of vector-observations is bounded (and the expected total number of observations required to terminate experimentation is uniformly in θ less than the total number required by the corresponding single-stage procedure of Bechhofer [1]).

In view of the desirable properties of the above procedures there appears to be little justification for using Tong's procedure.

REFERENCES

- [1] Bechhofer, R.E. (1954). A single-sample multiple decision procedure for ranking means of normal populations with known variances. Ann. Math. Statist., 25, 16-39.
- [2] Bechhofer, R.E., Kiefer, J., and Sobel, M. (1968). Sequential Identification and Ranking Procedures. University of Chicago Press, Chicago.
- [3] Chow, Y.S., Robbins, H., and Teicher, H. (1965). Moments of randomly stopped sums. Ann. Math. Statist., 36, 789-799.
- [4] Fabian, V. (1974). Note on Anderson's sequential procedures with triangular boundary. Ann. Statist. 2, 170-176.
- [5] Gibbons, J.D., Olkin, I., and Sobel, M. (1977). Selecting and Ordering Populations. Wiley, New York.
- [6] Paulson, E. (1964). A sequential procedure for selecting the population with the largest mean from k normal populations. Ann. Math. Statist., 35, 174-180.
- [7] Tamhane, A.C. and Bechhofer, R.E. (1977). A two-stage minimax procedure with screening for selecting the largest normal mean. Commun. Statist. - Theor. Meth., A6, 1003-1033.
- [8] Tamhane, A.C. and Bechhofer, R.E. (1979). A two-stage minimax procedure with screening for selecting the largest normal mean (II): an improved PCS lower bound and associated tables. Commun. Statist. - Theor. Meth., A8, 337-358.
- [9] Tong, Y.L. (1978). An adaptive solution to ranking and selection problems. Ann. Statist., 6, 658-672.

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

14 TR-480

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER No. 480 ✓	2. GOVT ACCESSION NO. AD-A099821	3. RECIPIENT'S CATALOG NUMBER (9)
4. TITLE (and Subtitle) Conditions Under Which $E(N_1) = \uparrow$ for Tong's Adaptive Solution To Ranking and Selection Problems.		5. TYPE OF REPORT & PERIOD COVERED Technical Report.
7. AUTHOR(s) Charles E. McCulloch		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS School of Operations Research and Industrial Engineering, College of Engineering, Cornell University, Ithaca NY 14853		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0036 N00014-75-C-0586
11. CONTROLLING OFFICE NAME AND ADDRESS (12) 10		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (11)
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Sponsoring Military Activities: US Army Research Office, P.O. Box 12211, Research Triangle Park, NC 27709, and Statistics and Probability Program, Office of Naval Research, Arlington, VA 22217		12. REPORT DATE September 1980
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited		13. NUMBER OF PAGES 4
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		14. SECURITY CLASS. (of this report) Unclassified
18. SUPPLEMENTARY NOTES		15. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ranking and selection procedures, adaptive sampling, sequential sampling, indifference-zone approach		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) ON REVERSE SIDE		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

409864

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Abstract

→ Tong [9] proposed an adaptive approach as an alternative to the classical indifference-zone formulation of the problems of ranking and selection. With a fixed pre-selected γ^* ($1/k < \gamma^* < 1$) his procedure calls for the termination of ^{going to} vector-at-a-time sampling when the estimated probability of a correct selection exceeds γ^* for the first time. The purpose of this note is to show that for the case of two normal populations with common known variance, the expected number of vector-observations required by Tong's procedure to terminate sampling approaches infinity as the two population means approach equality for $k \geq 3$ population means approach equality. Since in the typical ranking and selection setting it usually is assumed that the experimenter has no knowledge concerning the differences between the population means, the experimenter who uses Tong's procedure clearly does so at his own risk.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)